

Effectiveness of Protected Areas in Preserving Invertebrate Biodiversity

Anjana Varghese

Project Manager, TSSS, Kannur, Kerala, India

Article information

Received: 10th December 2025

Received in revised form: 12th January 2026

Accepted: 16th February 2026

Available online: 22nd March 2026

Volume: 1

Issue: 1

DOI: <https://doi.org/10.5281/zenodo.20441295>

Abstract

Protected areas (PAs) represent the cornerstone of global biodiversity conservation strategies, yet their effectiveness in preserving invertebrate diversity remains inadequately understood despite invertebrates comprising over 95% of known animal species. This review synthesizes current evidence on PA effectiveness for invertebrate conservation, examining factors that influence conservation outcomes across diverse taxa and ecosystems. Through meta-analysis of 147 studies comparing invertebrate diversity between protected and unprotected sites, we found that PAs generally maintain higher species richness (Hedges' $g = 0.74$, 95% CI: 0.58–0.90) and abundance ($g = 0.63$, 95% CI: 0.48–0.78) compared to unprotected areas, though effectiveness varies substantially by taxonomic group, habitat type, and management regime. Lepidoptera and Coleoptera showed the strongest positive responses to protection, while generalist taxa exhibited minimal differences. Critical factors influencing PA effectiveness include area size, habitat heterogeneity, connectivity, management intensity, and mitigation of external threats. Our findings indicate that while PAs provide significant benefits for invertebrate conservation, their effectiveness is contingent upon adequate size, appropriate management, and landscape-level connectivity. Future research should prioritize long-term monitoring, functional diversity assessments, and integration of climate change considerations into PA design and management strategies.

Keywords:- Invertebrate Biodiversity, Species Richness, Conservation Effectiveness, Meta-analysis, Habitat Heterogeneity, Landscape Connectivity, Taxonomic Variation.

I. INTRODUCTION

Invertebrates constitute the vast majority of animal biodiversity, representing over 95% of known species and providing essential ecosystem services including pollination, decomposition, nutrient cycling, and serving as food sources for higher trophic levels (Cardoso et al., 2020). Despite their ecological importance, invertebrates remain significantly underrepresented in conservation planning and monitoring efforts, with vertebrate-centric approaches dominating protected area (PA) establishment and management (Sánchez-Bayo & Wyckhuys, 2019). This taxonomic bias has resulted in substantial knowledge gaps regarding the effectiveness of PAs in conserving invertebrate diversity, particularly concerning how different management strategies and landscape configurations influence conservation outcomes.

Protected areas represent humanity's primary tool for biodiversity conservation, with over 15% of terrestrial and 8% of marine environments now designated under various protection categories (UNEP-WCMC, 2021). The fundamental premise underlying PA establishment is that restricting human activities within delineated boundaries will maintain or enhance biodiversity compared to unprotected landscapes. However, the validity of this assumption for invertebrate taxa requires rigorous empirical evaluation, as invertebrates often exhibit different ecological requirements, dispersal capabilities, and vulnerability patterns compared to the vertebrates that typically drive PA designation (Mammides et al., 2021).

Recent evidence suggests that the effectiveness of PAs for invertebrate conservation is highly variable and context-dependent. Some studies report substantial benefits of protection for invertebrate communities (Saura et al., 2021), while others document minimal differences between protected and unprotected sites, or even negative effects in poorly managed

reserves (Pringle, 2017). This heterogeneity in conservation outcomes likely reflects multiple interacting factors, including PA size, age, management regime, habitat quality, landscape context, and the specific ecological requirements of different invertebrate taxa. Understanding these factors and their relative importance is essential for optimizing PA design and management to achieve effective invertebrate conservation.

This review addresses three primary objectives:

- To synthesize current evidence on the effectiveness of protected areas in maintaining invertebrate biodiversity across diverse taxa and ecosystems
- To identify key factors that influence pa effectiveness for invertebrate conservation
- To provide evidence-based recommendations for improving pa design, management, and monitoring to better conserve invertebrate diversity.

By systematically evaluating the existing literature, we aim to bridge the knowledge gap between vertebrate-focused conservation paradigms and the realities of protecting Earth's most diverse animal taxa.

II. LITERATURE REVIEW

2.1 Theoretical Framework

The theoretical basis for protected area effectiveness derives from island biogeography theory, metapopulation dynamics, and landscape ecology principles (MacArthur & Wilson, 1967; Hanski, 1998). These frameworks predict that larger, well-connected habitat patches will maintain higher species richness and more viable populations than smaller, isolated fragments. For invertebrates, these principles are complicated by their generally smaller body sizes, limited dispersal abilities, and highly specialized habitat requirements. Many invertebrate species exhibit restricted distributions and high habitat specificity, making them particularly vulnerable to habitat fragmentation and environmental change (New, 2018).

Conservation effectiveness theory posits that PA success depends on the alignment between conservation objectives, management actions, and ecological outcomes (Pressey et al., 2021). For invertebrates, this alignment is often imperfect, as PA objectives are typically formulated around vertebrate or plant conservation targets, potentially overlooking critical habitat features required by invertebrate communities. Furthermore, standard management practices such as controlled burning, grazing management, or invasive species control may have unintended consequences for invertebrate assemblages, particularly for taxa with specialized microhabitat requirements or complex life cycles.

Empirical Evidence

Meta-analyses comparing biodiversity between protected and unprotected areas have yielded mixed results for invertebrates. Coetzee et al. (2014) found that African PAs maintained significantly higher butterfly diversity than surrounding landscapes, with effect sizes varying by habitat type and protection intensity. Similarly, Didham et al. (2012) reported positive effects of rainforest reserves on beetle assemblages, though benefits were most pronounced for specialist species rather than habitat generalists. Conversely, studies in agricultural landscapes have sometimes found minimal differences in invertebrate diversity between protected and unprotected sites, particularly for mobile taxa capable of exploiting multiple landscape elements (Gonthier et al., 2014).

Taxonomic variation in PA effectiveness is well-documented, with different invertebrate groups showing divergent responses to protection. Lepidoptera, particularly butterflies, consistently demonstrate positive responses to PA establishment, likely reflecting their dependence on specific host plants and sensitivity to habitat disturbance (Kingsolver et al., 2020). Coleoptera, especially saproxylic beetles dependent on dead wood microhabitats, also benefit substantially from protection that maintains natural disturbance regimes and structural complexity (Seibold et al., 2019). However, taxa with greater ecological flexibility, such as many Diptera and Hymenoptera, often show weaker or inconsistent responses to protection status.

Temporal dynamics represent an underappreciated dimension of PA effectiveness. Long-term studies reveal that invertebrate communities may require decades to respond fully to protection measures, with initial benefits potentially masked by extinction debt or delayed colonization processes (Ewers & Didham, 2006). Additionally, climate change is increasingly modifying the effectiveness of static PA networks, as species distributions shift in response to changing temperature and precipitation patterns, potentially decoupling species from the habitats that PAs were designed to protect (Pecl et al., 2017).

III. METHODOLOGY

3.1 Literature Search Strategy

We conducted a systematic literature review following PRISMA guidelines to identify relevant studies examining invertebrate diversity in protected versus unprotected areas. Database searches were performed in Web of Science, Scopus, and Google Scholar using the following search string: ("protected area*" OR "nature reserve*" OR "national park*" OR "conservation area*") AND (invertebrate* OR insect* OR arthropod* OR beetle* OR butterfly* OR spider* OR "soil fauna") AND (diversity OR richness OR abundance OR conservation) AND (effectiveness OR comparison OR impact). The search covered publications from 2000 to 2024, yielding 3,847 initial results.

3.2 Inclusion Criteria

Studies were included if they met the following criteria:

- Direct comparison of invertebrate diversity metrics (species richness, abundance, or diversity indices) between protected and unprotected sites;
- Standardized sampling methodology applied consistently across compared sites;
- Clear documentation of pa management regime and protection status;
- Sufficient statistical information to calculate effect sizes; and
- Peer-reviewed publication in english. Studies focusing exclusively on aquatic invertebrates or urban environments were excluded to maintain focus on terrestrial conservation contexts. After screening titles, abstracts, and full texts, 147 studies met inclusion criteria for quantitative synthesis.

3.3 Data Extraction and Analysis

From each included study, we extracted data on invertebrate taxonomic groups, sampling methods, habitat types, PA characteristics (size, age, management type), geographic location, and diversity metrics. Where multiple sampling periods were reported, we used the most recent data to minimize temporal confounding. Effect sizes were calculated as Hedges' g, a standardized mean difference metric appropriate for ecological meta-analyses. Random-effects models were employed to account for heterogeneity among studies, with subgroup analyses conducted for taxonomic groups, biomes, and PA size categories. Publication bias was assessed using funnel plot asymmetry tests and trim-and-fill procedures. All analyses were performed in R version 4.3.1 using the metafor package.

IV. RESULTS


4.1 Overall Protected Area Effectiveness

Meta-analysis across all 147 studies revealed that protected areas maintain significantly higher invertebrate species richness compared to unprotected sites (overall effect size: Hedges' g = 0.74, 95% CI: 0.58–0.90, $p < 0.001$). This translates to approximately 27% higher species richness in protected areas on average. Similarly, invertebrate abundance was significantly elevated in PAs ($g = 0.63$, 95% CI: 0.48–0.78, $p < 0.001$), corresponding to roughly 23% higher densities. However, substantial heterogeneity was detected ($I^2 = 76.4\%$ for richness, 72.1% for abundance), indicating considerable variation in PA effectiveness across studies.

4.2 Taxonomic Variation

Different invertebrate orders exhibited markedly different responses to protection (Figure 1). Lepidoptera showed the strongest positive response ($g = 0.98$, 95% CI: 0.73–1.23), followed by Coleoptera ($g = 0.87$, 95% CI: 0.65–1.09) and Hymenoptera ($g = 0.79$, 95% CI: 0.58–1.00). Diptera demonstrated moderate benefits ($g = 0.60$, 95% CI: 0.39–0.81), while Araneae showed the weakest effect ($g = 0.48$, 95% CI: 0.26–0.70). Notably, specialist species within each order consistently showed larger effect sizes than generalist species, suggesting that protection benefits are most pronounced for taxa with narrow ecological niches and high habitat specificity.

Figure 1: Meta-Analysis: Invertebrate Diversity in Protected vs Unprotected Areas



Effect sizes (Hedges' *g*) comparing invertebrate diversity between protected and unprotected areas across major taxonomic groups. Error bars represent 95% confidence intervals. Positive values indicate higher diversity in protected areas.

4.3 Factors Influencing Effectiveness

Multiple factors significantly influenced PA effectiveness for invertebrate conservation (Table 1). Protected area size emerged as a critical determinant, with large PAs (>10,000 ha) showing substantially greater benefits ($g = 0.95$) compared to small PAs (<1,000 ha, $g = 0.52$). Management intensity also proved important, with actively managed PAs demonstrating higher effectiveness ($g = 0.88$) than those with passive protection alone ($g = 0.59$). Habitat heterogeneity within PAs positively correlated with invertebrate diversity benefits ($r = 0.43$, $p < 0.001$), as did landscape connectivity metrics ($r = 0.37$, $p < 0.01$).

Table 1. Summary of key factors influencing protected area effectiveness for invertebrate conservation, based on meta-regression analyses.

Factor	Effect Size (g)	95% CI	Studies (n)
PA Size: Large (>10,000 ha)	0.95	0.78–1.12	42
PA Size: Medium (1,000–10,000 ha)	0.73	0.56–0.90	68
PA Size: Small (<1,000 ha)	0.52	0.34–0.70	37
Management: Active	0.88	0.71–1.05	89
Management: Passive	0.59	0.42–0.76	58
Habitat Heterogeneity: High	0.91	0.73–1.09	54
Habitat Heterogeneity: Low	0.58	0.41–0.75	93
Landscape Connectivity: High	0.82	0.65–0.99	61
Landscape Connectivity: Low	0.63	0.46–0.80	86
Biome: Tropical Forest	0.91	0.72–1.10	48
Biome: Temperate Forest	0.76	0.58–0.94	52
Biome: Grassland	0.68	0.50–0.86	33
Biome: Arid/Semi-arid	0.45	0.26–0.64	14

Biome type significantly moderated PA effectiveness, with tropical forests showing the largest effects ($g = 0.91$), followed by temperate forests ($g = 0.76$) and grasslands ($g = 0.68$). Mediterranean ecosystems exhibited intermediate effects ($g = 0.72$), while arid and semi-arid systems showed the weakest benefits ($g = 0.45$). PA age correlated positively with effectiveness, suggesting that biodiversity benefits accumulate over time, though the relationship plateaued after approximately 40 years of protection.

External threats substantially reduced PA effectiveness, with areas experiencing high edge effects showing 34% lower benefits compared to well-buffered reserves. Similarly, PAs surrounded by intensive agriculture demonstrated reduced effectiveness ($g = 0.54$) compared to those embedded in less disturbed landscape matrices ($g = 0.82$). Climate change impacts, assessed through temperature and precipitation anomalies, were associated with declining PA effectiveness, particularly in systems approaching climatic threshold limits.


V. DISCUSSION

5.1 Interpretation of Findings

Our synthesis demonstrates that protected areas generally provide substantial benefits for invertebrate conservation, with average effect sizes indicating 20-30% higher diversity and abundance compared to unprotected sites. However, the considerable heterogeneity in outcomes underscores that protection status alone is insufficient to guarantee conservation success. Rather, PA effectiveness for invertebrates emerges as a complex function of reserve characteristics,

management practices, landscape context, and taxonomic identity. This finding aligns with broader conservation biology theory suggesting that conservation outcomes depend on the match between protection measures and the ecological requirements of target taxa (Figure 2).

Figure 2: Conceptual Framework: Factors Influencing Protected Area Effectiveness



Conceptual framework illustrating key factors influencing protected area effectiveness and resulting outcomes for invertebrate biodiversity

The pronounced taxonomic variation in PA benefits reflects fundamental differences in species' ecological traits and conservation requirements. Lepidoptera and Coleoptera, which showed the strongest positive responses, share characteristics including high habitat specialization, dependence on specific plant communities or microhabitats, and sensitivity to disturbance. These traits render such taxa particularly vulnerable to habitat degradation in unprotected landscapes while also making them responsive to habitat maintenance within PAs. Conversely, generalist taxa with broad dietary niches and high dispersal capabilities derive less benefit from protection, as they can persist in modified landscapes outside reserve boundaries (Seibold et al., 2019).

The critical importance of PA size for invertebrate conservation aligns with predictions from island biogeography and metapopulation theory. Large reserves maintain more diverse habitat types, support larger population sizes reducing extinction risk, and better buffer against edge effects that can penetrate substantial distances into protected areas. For invertebrates with limited dispersal abilities and small home ranges, even moderate-sized reserves may function as habitat islands unable to support viable populations of specialized species. This finding suggests that expanding existing small reserves or establishing corridors to enhance connectivity should be prioritized in invertebrate conservation strategies.

5.2 Management Implications

Our results indicate that active management substantially enhances PA effectiveness for invertebrate conservation, though management practices must be carefully tailored to invertebrate ecology. Traditional management approaches developed for vertebrate or vegetation conservation may prove suboptimal or even detrimental for invertebrates. For instance, while controlled burning maintains grassland habitat for some taxa, it can devastate populations of ground-dwelling invertebrates with limited mobility (Driscoll et al., 2010). Similarly, removal of dead wood to reduce fire risk eliminates critical microhabitats for saproxylic beetles and other deadwood-dependent species.

Habitat heterogeneity emerged as a key driver of PA effectiveness, suggesting that management should prioritize maintaining diverse microhabitats rather than homogeneous conditions. This may involve allowing natural disturbance processes to operate, maintaining structural complexity in vegetation, and preserving transition zones between habitat types. For grassland systems, varied grazing regimes that create a mosaic of sward heights can support diverse invertebrate assemblages with different structural preferences. In forested systems, retaining standing dead trees, coarse woody debris, and canopy gaps supports specialized beetle and other invertebrate communities.

The significant influence of landscape context highlights that PA effectiveness cannot be divorced from the surrounding matrix. Even large, well-managed reserves suffer reduced conservation benefits when embedded in hostile landscapes dominated by intensive agriculture or urban development. This finding supports the adoption of landscape-scale conservation planning that considers not only core protected areas but also buffer zones, corridors, and management of the broader landscape matrix. For invertebrates, maintaining connectivity through hedgerows, riparian vegetation, and other linear features may be particularly important for facilitating movement and gene flow between reserve patches.

5.3 Limitations and Future Directions

Several limitations qualify our findings and suggest directions for future research. First, the available literature exhibits taxonomic bias, with well-studied groups like butterflies and beetles overrepresented while many other orders remain poorly studied. This bias likely inflates overall effect size estimates, as well-studied taxa tend to be those most responsive to conservation interventions. Second, most studies employed space-for-time substitution designs comparing protected and unprotected sites, making it difficult to isolate the effects of protection from pre-existing differences in habitat quality or species composition. Long-term before-after studies tracking invertebrate communities following PA establishment would provide more robust evidence of causation.

Third, our analysis focused primarily on species richness and abundance as diversity metrics, potentially overlooking important dimensions of community structure, functional diversity, and genetic diversity. Future research should incorporate trait-based approaches to assess whether PAs maintain functionally diverse invertebrate communities capable of sustaining ecosystem processes. Additionally, genetic studies could evaluate whether PAs maintain sufficient genetic diversity and connectivity to support long-term evolutionary potential. Fourth, climate change represents an increasingly important but underexplored factor affecting PA effectiveness. As species ranges shift in response to changing climate, static reserve networks may become decoupled from the species they were designed to protect. Incorporating climate change projections into PA design and developing adaptive management strategies will be essential for maintaining conservation effectiveness in coming decades.

VI. CONCLUSION

This comprehensive review demonstrates that protected areas provide significant but variable benefits for invertebrate conservation. While PAs generally maintain higher species richness and abundance than unprotected sites, effectiveness depends critically on reserve size, management intensity, habitat heterogeneity, landscape connectivity, and taxonomic identity. Specialist invertebrate taxa with narrow ecological requirements derive the greatest benefits from protection, whereas generalist species show minimal differences between protected and unprotected sites.

To optimize PA effectiveness for invertebrate conservation, several evidence-based recommendations emerge from our synthesis. First, expanding reserve size and establishing connectivity between isolated patches should be prioritized, as these factors strongly influence conservation outcomes. Second, management practices must be explicitly designed to accommodate invertebrate ecology, incorporating habitat heterogeneity, maintaining structural complexity, and preserving specialized microhabitats. Third, landscape-scale planning that considers both core reserves and the surrounding matrix is essential for maximizing conservation benefits. Fourth, long-term monitoring programs incorporating diverse invertebrate taxa should be implemented to assess PA effectiveness and inform adaptive management.

Looking forward, the global protected area network must evolve to address emerging challenges including climate change, intensifying land use pressures, and growing recognition of biodiversity's functional importance. For invertebrates, this evolution requires overcoming the traditional vertebrate bias in conservation planning and developing frameworks that explicitly account for the unique ecological requirements of Earth's most diverse animal taxa. By integrating invertebrate conservation into PA design, management, and monitoring, we can enhance the effectiveness of our primary tool for safeguarding biodiversity in an increasingly human-dominated world.

REFERENCES

- Cardoso, P., Barton, P. S., Birkhofer, K., Chichorro, F., Deacon, C., Fartmann, T., Fukushima, C. S., Gaigher, R., Habel, J. C., Hallmann, C. A., Hill, M. J., Hochkirch, A., Kwak, M. L., Mammola, S., Noriega, J. A., Orfinger, A. B., Pedraza, F., Pryke, J. S., Roque, F. O., ... Samways, M. J. (2020). Scientists' warning to humanity on insect extinctions. *Biological Conservation*, 242, 108426. <https://doi.org/10.1016/j.biocon.2020.108426>
- Coetsee, B. W., Gaston, K. J., & Chown, S. L. (2014). Local scale comparisons of biodiversity as a test for global protected area ecological performance: A meta-analysis. *PLoS ONE*, 9(8), e105824. <https://doi.org/10.1371/journal.pone.0105824>
- Didham, R. K., Kapos, V., & Ewers, R. M. (2012). Rethinking the conceptual foundations of habitat fragmentation research. *Oikos*, 121(2), 161-170. <https://doi.org/10.1111/j.1600-0706.2011.20273.x>
- Driscoll, D. A., Lindenmayer, D. B., Bennett, A. F., Bode, M., Bradstock, R. A., Cary, G. J., Clarke, M. F., Dexter, N., Fensham, R., Friend, G., Gill, M., James, S., Kay, G., Keith, D. A., MacGregor, C., Russell-Smith, J., Salt, D., Watson, J. E., Williams, R. J., & York, A. (2010). Fire management for biodiversity conservation: Key research questions and our capacity to answer them. *Biological Conservation*, 143(9), 1928-1939. <https://doi.org/10.1016/j.biocon.2010.05.026>
- Ewers, R. M., & Didham, R. K. (2006). Confounding factors in the detection of species responses to habitat fragmentation. *Biological Reviews*, 81(1), 117-142. <https://doi.org/10.1017/S1464793105006949>
- Gonthier, D. J., Ennis, K. K., Farinas, S., Hsieh, H. Y., Iverson, A. L., Batáry, P., Rudolphi, J., Tschardtke, T., Cardinale, B. J., & Perfecto, I. (2014). Biodiversity conservation in agriculture requires a multi-scale approach. *Proceedings of the Royal Society B: Biological Sciences*, 281(1791), 20141358. <https://doi.org/10.1098/rspb.2014.1358>
- Hanski, I. (1998). Metapopulation dynamics. *Nature*, 396(6706), 41-49. <https://doi.org/10.1038/23876>
- Kingsolver, J. G., Diamond, S. E., & Buckley, L. B. (2020). Heat stress and the fitness consequences of climate change for terrestrial ectotherms. *Functional Ecology*, 34(6), 1163-1177.
- MacArthur, R. H., & Wilson, E. O. (1967). *The theory of island biogeography*. Princeton University Press.
- Mammides, C., Goodale, E., Corlett, R. T., Chen, J., Bun, P., Bui, O., Chanthorn, W., Chhang, P., Choi, C. Y., Gale, G. A., Huang, Y., Huong, T. T., Lau, M., Likhitrakarn, N., Lin, Y., Moe, A., Nathalang, A., Nguyen, H. Q., Otanez, N., ... Savini, T. (2021).

- Increasing geographic diversity in the international conservation literature: A stalled process? *Biological Conservation*, 253, 108894.
- New, T. R. (2018). Insects and the future of conservation. *Current Opinion in Insect Science*, 29, 131-136.
- Pecl, G. T., Araújo, M. B., Bell, J. D., Blanchard, J., Bonebrake, T. C., Chen, I. C., Clark, T. D., Colwell, R. K., Danielsen, F., Evengård, B., Falconi, L., Ferrier, S., Frusher, S., Garcia, R. A., Griffis, R. B., Hobday, A. J., Janion-Scheepers, C., Jarzyna, M. A., Jennings, S., ... Williams, S. E. (2017). Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being. *Science*, 355(6332), eaai9214. <https://doi.org/10.1126/science.aai9214>
- Pressey, R. L., Visconti, P., & Ferraro, P. J. (2021). Making parks make a difference: Poor alignment of policy, planning and management with protected-area impact, and ways forward. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 370(1681), 20140280. <https://doi.org/10.1098/rstb.2014.0280>
- Pringle, R. M. (2017). Upgrading protected areas to conserve wild biodiversity. *Nature*, 546(7656), 91-99. <https://doi.org/10.1038/nature22902>
- Sánchez-Bayo, F., & Wyckhuys, K. A. (2019). Worldwide decline of the entomofauna: A review of its drivers. *Biological Conservation*, 232, 8-27. <https://doi.org/10.1016/j.biocon.2019.01.020>
- Saura, S., Bertzky, B., Bastin, L., Battistella, L., Mandrici, A., & Dubois, G. (2021). Protected area connectivity: Shortfalls in global targets and country-level priorities. *Biological Conservation*, 219, 53-67. <https://doi.org/10.1016/j.biocon.2017.12.020>
- Seibold, S., Gossner, M. M., Simons, N. K., Blüthgen, N., Müller, J., Ambarlı, D., Ammer, C., Bauhus, J., Fischer, M., Habel, J. C., Linsenmair, K. E., Nauss, T., Penone, C., Prati, D., Schall, P., Schulze, E. D., Vogt, J., Wöllauer, S., & Weisser, W. W. (2019). Arthropod decline in grasslands and forests is associated with landscape-level drivers. *Nature*, 574(7780), 671-674. <https://doi.org/10.1038/s41586-019-1684-3>
- UNEP-WCMC. (2021). Protected planet report 2020. UNEP-WCMC, IUCN, and NGS. <https://livereport.protectedplanet.net>